

2017 Location Sheet

Location: Farmland

**Corn**

Population: 35,000

Previous crop: Soybean

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Blount - Pewamo

**Fertility Rates**

N: 216 lbs/a

P: None

K: None

**Herbicides:** Bucaneer 5 Xtra  
Choice Weathermaster  
Charger Max ATZ

**Insecticide:** None

**Tillage:**

Fall: chisel  
\_\_\_\_\_

Spring: field cultivator  
\_\_\_\_\_  
\_\_\_\_\_

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0.00

30yr. Average 23.21

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Blount - Pewamo

**Fertility Rates**

N: None

P: None

K: None

**Herbicides:** Dual II Magnum  
Canopy XL  
Bucaneer 5 Xtra  
Choice Weathermaster

**Insecticide:** None

**Tillage:**

Fall: disc  
\_\_\_\_\_

Spring: field cultivator X 2  
\_\_\_\_\_  
\_\_\_\_\_

2017 Location Sheet

Location: Butlerville

**Corn**

Population: 35,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Avonburg Silt loam

**Fertility Rates**

N: 200# N

P: 250# DAP

K: 400# Potash)

Herbicides: Halex GT

Fungicide: None

Insecticide: None

**Tillage:**

Fall: None

Spring: Disc 2X

Soil Finisher

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0.00

30yr. Average 22.97

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Avonburg/Cobbsfork

**Fertility Rates**

N: \_\_\_\_\_

P: 250# DAP

K: 277# Potash

Herbicides: \_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_

Spring: Disc 2X

Soil Finisher

2017 Location Sheet

Location: Vincennes

**Corn**

Population: 28,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Paton Silt Loam

**Fertility Rates**

N: 170#

P: \_\_\_\_\_

K: 72#

Herbicides: Harness Extra

Round Up Power Max

\_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Chisel

\_\_\_\_\_

Spring: Field cultivated

Disked twice

\_\_\_\_\_

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0

30yr. Average 24.75

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Henshaw Slit Loam

**Fertility Rates**

N: 23#

P: \_\_\_\_\_

K: 72#

Herbicides: Scepter

Dual

Round-up Power Max

\_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Chisel

\_\_\_\_\_

Spring: Field cultivated

Disked twice

\_\_\_\_\_

2017 Location Sheet

Location: Wanatah

**Corn**

Population:35,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Sebowa Loam

**Fertility Rates**

N: 30#pre/180#sideUAN

P: 80 lb

K: 100 lb

Herbicides: \_\_\_\_\_

Fultime ppi

Calisto Xtra

Fungicide: \_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Chisel Plow

Spring: Field Cultivate 1X

\_\_\_\_\_

\_\_\_\_\_

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0.00

30yr. Average 22.87

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Sebowa Loam

**Fertility Rates**

N: \_\_\_\_\_

P: \_\_\_\_\_

K: \_\_\_\_\_

Herbicides: Canopy DF ppi

FlexStar post

First Rate

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Chisel Plow

Spring: Disk 1X

Field Cultivate 1X

\_\_\_\_\_

2017 Location Sheet

Location: Columbia City

**Corn**

Population: 35,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Rawson Sandy Loam

**Fertility Rates**

N: 188 lb N/a (planned)

P: 66 lb P2O5/a

K: 70 lb K2O/a

Herbicides: Atrazine

Dual

2,4-D

Fungicide: none yet

Insecticide: Warhawk

**Tillage:**

Fall: \_\_\_\_\_

Spring: vertical tillage tool x 2

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0

30yr. Average 21.10

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Rawson Sandy Loam

**Fertility Rates**

N: 0

P: 60 lb P2O5/a

K: 125 lb K2O/a

Herbicides: Sonic

glyphosate

\_\_\_\_\_

none yet

Insecticide: none yet

**Tillage:**

Fall: deep ripper

Spring: vertical tillage tool x2

2017 Location Sheet

Location: Shelburn

**Corn**

Population: 33,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Iva Silt Loam

**Fertility Rates**

N: 200#

P: \_\_\_\_\_

K: \_\_\_\_\_

Herbicides: 1/2 rate Degree

Atrazine

Roundup

Fungicide: \_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_

\_\_\_\_\_

Spring: Disk

Field cultivator

\_\_\_\_\_

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0.00

30yr. Average 23.76

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Iva Silt Loam

**Fertility Rates**

N: \_\_\_\_\_

P: \_\_\_\_\_

K: \_\_\_\_\_

Herbicides: Roundup

\_\_\_\_\_

\_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_

\_\_\_\_\_

Spring: Chisel

Field cultivator

\_\_\_\_\_

**2017 Location Sheet**

**Location:** Rensselaer

**Corn**

Population: 36,000

**Previous crop:** Corn

**Planting Date:** \_\_\_\_\_

**Harvest Date:** \_\_\_\_\_

**Soil Type:** Zadog-Grandby Complex

**Fertility Rates**

N: 200#

P: 83#

K: 180#

**Herbicides:** Durango, Keystone

Atrazine

Laudis

**Fungicide:** \_\_\_\_\_

**Insecticide:** \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_

Spring: field cultivated 2X

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

Total **0.00**

30yr. Average 22.52

2017 Location Sheet

Location: Milford

**Corn**

Population: 32,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Croiser Loam

**Fertility Rates**

N: 145 lbs

P: variable rate

K: variable rate

Herbicides: atrazine

roundup

Fungicide: \_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Chisel

\_\_\_\_\_

Spring: Disk

Field cultivated

\_\_\_\_\_

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

Total **0.00**

30yr. Average 21.07

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Croiser Loam

**Fertility Rates**

N: Dairy manure

P: variable rate

K: variable rate

Herbicides: Burndown

roundup

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_

\_\_\_\_\_

Spring: Tubro till x2

\_\_\_\_\_

\_\_\_\_\_



## 2017 Location Sheet

**Location:** Washington

### Corn

Population: 36,000

**Previous crop:** Corn

**Planting Date:** \_\_\_\_\_

**Harvest Date:** \_\_\_\_\_

**Soil Type:** Iva Silt Loam

### Fertility Rates

**N:** 225 lbs

**P:** 3tons turkey manure

**K:** 120 lbs

**Herbicides:** Roundup weathermax

**Fungicide:** yes

**Insecticide:** \_\_\_\_\_

### Tillage:

**Fall:** Chisel plow

**Spring:** JD finishing tool

### Monthly Rainfall:

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

Total **0.00**

30yr. Average **23.75**

2017 Location Sheet

Location: Tipton

**Corn**

Population:35,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: \_\_\_\_\_

**Fertility Rates**

N: \_\_\_\_\_  
P: \_\_\_\_\_  
K: \_\_\_\_\_

Herbicides: \_\_\_\_\_  
\_\_\_\_\_

Fungicide: \_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_  
\_\_\_\_\_

Spring: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**Monthly Rainfall:**

April	_____
May	_____
June	_____
July	_____
August	_____
September	_____
Total	<b>0.00</b>
30yr. Average	<u>22.72</u>

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: \_\_\_\_\_

**Fertility Rates**

N: \_\_\_\_\_  
P: \_\_\_\_\_  
K: \_\_\_\_\_

Herbicides: \_\_\_\_\_  
\_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: \_\_\_\_\_  
\_\_\_\_\_

Spring: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

**Monthly Rainfall:**

April	_____
May	_____
June	_____
July	_____
August	_____
September	_____
Total	<b>0.00</b>
30yr. Average	<u>22.97</u>

**2017 Location Sheet**

**Location:** Rushville

**Corn**

Population: 34,000

**Previous crop:** tomatoes

**Planting Date:** \_\_\_\_\_

**Harvest Date:** \_\_\_\_\_

**Soil Type:** Xenia silt loam

**Fertility Rates**

N: 220lbs

P: 0

K: 0

**Herbicides:** Roundup & Prowl

**Fungicide:** \_\_\_\_\_

**Insecticide:** \_\_\_\_\_

**Tillage:**

Fall: Ripped

Spring: Field cultivated 2X

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

Total **0.00**

30yr. Average 25.02

2017 Location Sheet

Location: W. Lafayette

**Corn**

Population: 35,000

Previous crop: Soybeans

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Chalmers silty clay loam

**Fertility Rates**

N: 180

P: 104

K: 180

Herbicides: Bicep II

Callisto

Fungicide: \_\_\_\_\_

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Ripped

Spring: Field Cultivator X2

**Monthly Rainfall:**

April \_\_\_\_\_

May \_\_\_\_\_

June \_\_\_\_\_

July \_\_\_\_\_

August \_\_\_\_\_

September \_\_\_\_\_

**Total** 0.00

30yr. Average 22.11

**Soybeans**

Previous crop: Corn

Planting Date: \_\_\_\_\_

Harvest Date: \_\_\_\_\_

Soil Type: Chalmers silty clay loam

**Fertility Rates**

N: \_\_\_\_\_

P: \_\_\_\_\_

K: \_\_\_\_\_

Herbicides: Duel

Septor

Roundup

Flexstar

Insecticide: \_\_\_\_\_

**Tillage:**

Fall: Chisel

Spring: Field Cultivator X2