

Herbicides for Cole Crops and Leafy Greens¹

Product (REI/PHI)	Common Name	Timing and Application Location Relative to Crop					Incorporated	Timing Relative to Weeds		Weed Groups Controlled			Crops				
		Before seeding	After seeding before emergence	Before transplanting	Post emergence — between rows only	Postemergence		Preemergence	Postemergence	Annual grasses	Small-seeded broadleaves	Broadleaves	Broccoli	Brussels sprouts	Cabbage	Cauliflower	Brassica greens
Aim EC (12h/-)	carfentrazone				X			X		X		X		X		X	
Command 3ME* (12h/45d)	clomazone	X	X	X				X	X	X		X		X			
Dacthal* (12h/-)	DCPA	X	X	X				X				X		X		X	
Devrinol 50DF* (12h/-)	napropamide	X	X	X			Yes	X	X	X		X		X			
Dual Magnum* (24h/21d)	s-metolachlor		X					X				X				X	
Goal 2XL* (12h/-)	oxyfluorfen			X				X	X	X		X		X			
Gramoxone* (12h to 24h/-)	paraquat	X	X	X					X	X		X		X		X	
Poast* (12h/14-30d)	sethoxydim	X	X	X					X			X		X		X	
Prefar 4E* (12h/-)	bensulide	X	X	X				X				X		X		X	
Roundup®, others (12h/14d)	glyphosate	X	X	X	X				X	X		X		X		X	
Select Max®, others (12h/14-30d)	clethodim	X	X	X					X	X		X		X		X	
Spartan DF* (12h/1d)	sulfentrazone			X	X			X				X					
Spartan Charge* (12h/-)	sulfentrazone plus carfentrazone			X	X			X		X							
Stinger®	clopyralid	X	X	X	X				X					X		X	
Treflan®, others (12h/-)	trifluralin	X		X			Yes	X				X		X		X	

¹For effectiveness against specific weeds, see Relative Effectiveness of Herbicides for Vegetable Crops (page 68), and read label. This table does not include all label information. Be sure to read and follow all instructions and precautions on the herbicide label. Herbicides can cause serious crop injury and yield loss if not used properly.