

August 1, 2018

Sulfur deficiency in alfalfa

Jim Camberato¹ and Keith Johnson

Agronomy Department, Purdue University, West Lafayette, IN

Sulfur (S) deficiency has recently been reported in a number of alfalfa fields in southwestern Wisconsin according to Carrie Laboski at the Univ. of Wisconsin (**see ref. 1**). Quite a few sulfur deficiencies in alfalfa and corn have been reported since 2005 in northeast Iowa by John Sawyer and Brian Lang (**see ref. 2, 3**). In Indiana we have documented S deficiency in wheat, corn, and soybean in recent years and are conducting some research on S deficiency as well (**see ref. 4**).

Factors Affecting Sulfur Deficiency

Sulfur deficiency of alfalfa and other crops may be becoming more prevalent because less S is deposited from the atmosphere to the soil due to reductions in power plant S emissions (**Fig. 1**). Higher yields result in greater crop S removal from the field - about 5-7 lb for each ton of alfalfa hay. Additionally, less incidental S applications in fertilizers and pesticides may contribute to more S deficiency. In the case of row crops, increases in no-till, early planting, and heavy residue from high yields have also been implicated in contributing to S deficiency.




Fig. 1. The amount of sulfur deposited on the land is much less in 2015 than in 2001. Red colors indicate high deposition and blue low deposition. Data from: <http://nadp.slh.wisc.edu/committees/tdep/tdepmaps/preview.aspx>. (URL accessed Aug. 2018).

¹For more information, contact J. Camberato (765-496-9338, jcambera@purdue.edu), or K. Johnson (765-494-4800, johnsonk@purdue.edu).

Soil Factors Resulting in Sulfur Deficiency

The main source of S in most soils is organic S. Each percent organic matter in the plow layer contains about 100 pounds of sulfur per acre. Organic S must be mineralized to sulfate-S ($\text{SO}_4\text{-S}$) to be taken up by crop plants, in much the same way that organic N is made available to crop plants. Therefore the lower the organic matter content of the soil the more likely S deficiency is to occur.

Since mineralization is a process carried out by living microorganisms, soil temperature and moisture largely determine when and how much of the organic S is made available to the crop. Cold and excessively wet or dry conditions reduce microbial activity and reduce S availability from soil organic matter and crop residues. Thus, alfalfa is more likely to be S deficient in the early spring before soil temperatures warm substantially.

Sulfate-S is relatively mobile in most soils (similar to nitrate) because it has a double negative charge and is repelled by the negative charge of the soil, unlike nutrients such as potassium, calcium, or magnesium. Although $\text{SO}_4\text{-S}$ can bind to iron and aluminum in the soil, these elements are much more likely to bind phosphate at the exclusion of $\text{SO}_4\text{-S}$ and as a result $\text{SO}_4\text{-S}$ is easily leached from soils, especially sandy soils.

At the field level the occurrence of S deficiency may be highly variable since soil S availability varies considerably with soil organic matter and texture. Sulfur deficiency is often seen in sandier, lower organic matter, and higher elevation areas of a field while lower lying, higher organic matter, and heavier textured areas have sufficient S.

Soil testing methods measure the $\text{SO}_4\text{-S}$ form of S. Unfortunately soil testing is not particularly useful for predicting S deficiency because it does not take into account the organic S component that might become available to the crop. The $\text{SO}_4\text{-S}$ component that is actually measured may also be leached from the soil between the time of sampling and the time of crop need.

Identifying Sulfur Deficiency in Alfalfa

At the individual plant level, S-deficient alfalfa will have a uniformly yellow appearance because S is not readily translocated from older to younger plant parts. Visually S deficiency may not be easy to distinguish from other nutrient deficiencies or leafhopper damage without close inspection, experience, and/or tissue testing.

The best way to identify a S deficiency is by tissue sampling. Collect the upper 6 inches of stem and leaf tissue from 40-50 plants from the area suspected of deficiency and a healthy area of the field for comparison. If samples are contaminated by soil they can be rinsed quickly in cold distilled water, but do not overdo it because some nutrients, especially potassium, may be leached out of the tissue. Wet samples should be air-dried before shipping to the laboratory in paper bags. Tissue S greater than about 0.25% on a dry matter basis indicates sufficiency.

Correcting Sulfur Deficiency in Alfalfa

If S deficiency is identified, an application rate of 20 to 30 pounds of SO₄-S per year is recommended to alleviate S deficiency in alfalfa based on the most recent research conducted in Iowa (**see ref. 3**). Although some carryover of S may occur in silt loam soils with deep rooted alfalfa crops it may be necessary to make applications of S every year on sandy soils, particularly if irrigated and high yielding.

Fertilizer Materials

The composition of several fertilizer sources of S are listed in **Table 1**. Potassium sulfate and sulfate-of-potash-magnesia (sul-po-mag or K-mag) are often good choices because alfalfa removes large amounts of potassium from the soil and potassium fertilization is often required. Either of these sources can be blended with muriate of potash (0-0-60) to provide an economical source of both potassium and S. The inclusion of magnesium in sul-po-mag may be an extra benefit compared to potassium sulfate if soil magnesium levels are low.

Ammonium sulfate and gypsum are also potential sources of S for alfalfa. Although there is no benefit to applying ammonium-nitrogen to the alfalfa, there are not likely to be any negative effects either with the rate that would be used to correct the S deficiency. Naturally-occurring mined gypsum and several by-product sources of gypsum can be applied to provide S as well. Unless pelletized, however, higher than necessary rates of S will be applied with gypsum which is difficult to spread at rates less than 500 to 1000 pounds per acre (85 to 170 pounds of S per acre assuming 17% S). If carryover of S occurs, the S will be utilized in later years. However, in sandy soils, where leaching is a problem, the benefit in future years may be minimal.

Table 1. Sulfur-containing fertilizers and their approximate composition. (**see ref. 5**). The actual nutrient concentration of a fertilizer should be expressed on the guarantee.

Fertilizer	%N	%K ₂ O	%S	%Mg	%Ca
Ammonium sulfate	21	0	24	0	0
Ammonium thiosulfate	12	0	26	0	0
Elemental sulfur	0	0	>90	0	0
Gypsum (calcium sulfate)	0	0	19	0	23
Potassium magnesium sulfate	0	22	23	11	0
Potassium sulfate	0	50	18	0	0

Effects of Sulfur Containing Fertilizers on Soil pH

Some of the S-containing fertilizers mentioned above affect soil pH. Elemental S, ammonium sulfate and ammonium thiosulfate generate acidity and reduce soil pH. No acidity arises from potassium, magnesium, or calcium sulfates.

References used in this article:

- (1) Laboski, Carrie. 2012. Sulfur fertilizer price comparison for alfalfa. Univ. Wisconsin Crop Manager for April 19, 2012. Available at: http://ipcm.wisc.edu/blog/2012/04/2012-sulfur-fertilizer-price-comparison-for-alfalfa/?utm_source=rss&utm_medium=rss&utm_campaign=2012-sulfur-fertilizer-price-comparison-for-alfalfa. (URL accessed August 2018).
- (2) Sawyer, John, Brian Lang, and D. Barker. 2011. Sulfur emerges as a nutritional issue in Iowa alfalfa production. Available at: http://www.agronext.iastate.edu/soilfertility/info/2006LangICMConfProcBL_11-16-06.pdf. (URL accessed August 2018).
- (3) Sawyer, John, Brian Lang, and Daniel Barker. 2011. Sulfur emerges as a nutritional issue in Iowa alfalfa production. Better Crops 95(2):6-7. Available at: http://www.agronext.iastate.edu/soilfertility/info/SulfurEmerges_Alfalfa_IPNIBetterCropsArticle.pdf. (URL accessed August 2018).
- (4) Camberato, Jim and Shaun Casteel. 2017. Sulfur deficiency. Purdue University Department of Agronomy, Soil Fertility Update. Available at: <https://ag.purdue.edu/agry/Documents/Sulfur%20deficiency%202017.pdf>. (URL accessed August 2018).
- (5) Crozier, C., D. Osmond, et al. 2017 North Carolina Agricultural Chemicals Manual - Fertilizer Use. <https://content.ces.ncsu.edu/north-carolina-agricultural-chemicals-manual/fertilizer-use>. (URL accessed August 2018).

© Purdue University, 2018

It is the policy of the Purdue University Cooperative Extension Service that all persons have equal opportunity and access to its educational programs, services, activities, and facilities without regard to race, religion, color, sex, age, national origin or ancestry, marital status, parental status, sexual orientation, disability or status as a veteran. Purdue University is an Affirmative Action institution. This material may be available in alternative formats.