

2017 Evaluation of Hybrid Bell Pepper Varieties for High Tunnel Production in Kansas

Kimberly L. Oxley and Cary L. Rivard, Kansas State University
Department of Horticulture and Natural Resources, Manhattan, KS koxley@ksu.edu

Introduction

High tunnel (hoop house) production of vegetables has become quite common in Kansas. High tunnels protect crops from harsh environmental conditions such as wind and storm damage. In addition to protection, determinate tomato crop requirements (planting date, soil temperature, crop height, etc.), high tunnels provide an excellent system for bell pepper production. Therefore, this method protects the crop, lengthens the growing season, and increases profitability for the grower as a result (Carey et al, 2009).

Bell peppers in Kansas are a valuable crop that is sold through farmers markets, CSA's, on-farm, wholesale and to restaurant. Results from a grower survey conducted in 2014 by the Kansas Rural center showed peppers were ranked fourth in popularity as one of the most common crops grown in high tunnels (Phelps, 2014). Similarly, peppers also ranked as the fifth most popular crop grown in high tunnels in the central United States (Knewton et al, 2010).

The goal of this study was to investigate the performance of ten hybrid bell pepper (green to red) varieties for fresh-market production in high tunnels. Ten commercially available varieties were tested and total yields ranged from 10.9 to 16.9 lb of fruit per plant. The three varieties with the highest marketable fruit number and weight per plant were 'Galileo,' 'Declaration,' and 'Archimedes.'

Materials and Methods

The trial was conducted at the Olathe Horticulture Research and Extension Center located approximately 30 miles southwest of Kansas City. Transplants were grown in soilless potting media using 50-cell propagation trays. Seeds were sown on 20 February 2017 and transplanted to 50-cell trays on 11 March. Transplants were set on 2 May in one bay of a multi-bay high tunnel (96' x 200' Haygrove Multi-Bay High Tunnel). The trial was planted into four rows with each row consisting of one replication. Each plot had five plants, and in-row spacing was 18", which is typical of commercial pepper production. Plastic mulch and drip irrigation were employed, and the stake-and-weave method was utilized as needed to trellis the plants vertically. Granular fertilizer was applied on 25 March at a rate of 50 lb N/acre. Weekly tissue samples were used to identify when fertigation was needed throughout the season. Fertigation was carried out at a rate of 10 lb nitrogen/acre and 5 lb/potassium/acre on 4 August. Harvesting occurred from 3 June through 9 October. During the last harvest, all fruit larger than 5 cm were picked. The fruit were graded for marketability, and fruit number and weight were recorded. Average fruit size and percent marketability were determined and are presented below. The data set was analyzed using an ANOVA (PlotIt, Scientific Programming Enterprises, Haslett, MI), and a mean separation test was carried out by using an F-protected least significant difference (LSD) test. A separate

analysis was carried out for each observation, and the results of the LSD test are shown where statistically significant treatment effects occurred.

Results and Discussion

In our trial, 'Declaration' yielded the highest marketable fruit number at 38.5 per plant and was not significantly different from 'Galileo', 'Archimedes', 'Vanguard', 'Red Knight', 'Karisma', or 'Intruder.' 'Archimedes', 'Intruder', and 'Declaration' had the highest percent marketability (by number) at 86.3%, 85.4%, and 84.9% respectively. 'Declaration' and 'Galileo' had similar marketable fruit sizes at approximately 0.38 lb. These varieties were not significantly different from 'Vanguard,' 'Karisma' or 'Archimedes' in regards to average marketable fruit size.

'Declaration' resulted in the highest marketable weight at 14.9 lb per plant and was not significantly different from 'Galileo', 'Archimedes', 'Vanguard', or 'Karisma.' 'Currier', 'Archimedes', 'Intruder' had the highest percent marketability (by weight) at 90.2%, 88.4%, and 88.1% respectively. Since 2013, 'Karisma', 'Declaration', and 'Archimedes' have consistently produced high marketable fruit number and weight per plant (data not shown), although 'Karisma' did not perform as well in 2017.

Our observations showed three flushes this season (data not shown) during harvests conducted on 20 July, 23 August, and 11 September. During the early peak harvest on 20 July, 'Vanguard,' 'Galileo,' and 'Vanguard' produced the highest total fruit number. On 23 August, 'Intruder,' 'Red Knight' and 'Declaration' produced the highest total fruit number. The harvest that occurred on 11 September was the largest and 'Archimedes,' 'Declaration' and 'Alliance' provided the highest total fruit number. The timing of market demand may determine which variety would be ideal for particular growers. If a grower is targeting an earlier season customer base, then 'Vanguard' and 'Galileo' may work best. However, if market demand is later in the season, then 'Declaration' and 'Archimedes' may be better-suited. A good mix would be ideal for growers who wants to have a consistent season-long supply.

'Karisma' has been trialed every year since 2013. In this trial, 'Karisma' was more consistent week-to-week throughout the duration of the season than any of the other varieties tested. It had a high early yield flush, but produced consistently for the remainder of the season with fewer fluctuations in marketable yield than other varieties (data not shown).

A large portion (>80%) of the fruit quality problems seen in this trial were the result of blossom end rot (BER). Though the culled fruit was not specifically graded for this issue, the effects observed in this study were most likely the result of a lower incidence of BER.

Table 1. Marketable and total per plant fruit yield of green pepper varieties grown in a three season high tunnel in Olathe, Kansas.

Variety	Marketable				Total			
	Number		Wt (lbs)		Number		Wt (lbs)	
Declaration	38.5	c	14.9	e	45.3	c	16.9	e
Galileo	38.3	c	14.4	de	45.4	c	16.7	e
Archimedes	37.9	c	13.3	cde	44.0	bc	15.0	cde
Vanguard	34.6	bc	13.1	cde	43.7	bc	15.7	de
Red Knight	34.0	bc	11.7	abcd	41.0	abc	13.7	abcd
Karisma	32.8	abc	12.2	bcde	38.8	abc	14.2	bcde
Intruder	32.2	abc	11.4	abc	37.3	ab	12.8	abc
Alliance	30.3	ab	10.7	abc	37.0	ab	12.5	abc
Currier	28.2	ab	9.9	ab	33.4	a	10.9	a
Bayonet	26.2	a	9.2	a	36.6	ab	11.9	ab
LSD _(0.05)	7.0355		2.8252		7.7760		2.9305	

Table 2. Mean pepper fruit size (lbs) and marketability of green pepper varieties grown in a three season high tunnel in Olathe, Kansas.

Variety	Average Fruit Size (lbs)				Percent Marketability			
	Marketable		Total		Number		Weight	
Declaration	0.38	b	0.37	c	84.9	b	87.8	bc
Galileo	0.38	ab	0.37	c	84.2	b	85.7	bc
Vanguard	0.37	ab	0.36	abc	79.0	ab	82.5	ab
Karisma	0.37	ab	0.36	bc	84.5	b	85.6	bc
Archimedes	0.35	ab	0.34	abc	86.3	b	88.4	bc
Intruder	0.35	a	0.34	abc	85.4	b	88.1	bc
Alliance	0.35	a	0.34	abc	81.6	b	84.4	bc
Currier	0.35	a	0.32	a	84.4	b	90.2	c
Bayonet	0.35	a	0.33	ab	71.8	a	75.6	a
Red Knight	0.34	a	0.34	abc	82.9	b	85.2	bc
LSD _(0.05)	0.0336		0.0390		9.4747		7.7531	

Acknowledgements

We would like to sincerely thank the Kansas Vegetable Growers Association for support of this project. Seeds were donated by Harris Moran, Seedway, and Johnny's Selected Seeds. Technical support provided by Paul Andersen, Zachary Callaghan, Zac Hoppenstedt, David Loewen, Tracy

Oelschlaeger, Connor Oxley, and Madison Tunnell. We also thank the Olathe Horticulture Research and Extension Center for assistance with this project.

Literature Cited

Carey, E.E., L. Jett, W.J. Lamont, T.T. Nennich, M.D. Orzolek, and K.A. Williams. 2009. Horticultural crop production in high tunnels in the United States: A snapshot. HortTechnology 19:37–43.

Knewton, S.J.B, Cary, E.E., Kirkham, M.B. 2010. Management Practices of Growers Using High Tunnels in the Central Great Plains of the United States. HortTechnology 20:639-645.

Phelps, Dan, November 2014. Growing Under Cover: A Guide to Polyunnel Options for Kansas Growers, a Kansas Rural Center publication, Topeka, KS.

U.S. Department of Agriculture. 2014. Census of Horticultural Specialties. Volume 3 Special Studies Part 3. State Data. Table 14. 2015 December.
agcensus.usda.gov/Publications/Census_of_Horticulture_Specialties

Seed Sources

Harris Moran - HM

Seedway – SW/SDW

Johnny’s Selected Seeds - JS

Contact Info

Kimberly Oxley

Research Associate

Kansas State University

35230 W 135th St

Olathe, KS 66061

koxley@ksu.edu

913-856-2335 x121